



Miskolc Mathematical Notes
Vol. 15 (2014), No 2, pp. 711-716

HU e-ISSN 1787-2413
DOI: 10.18514/MMN.2014.640

An existence theorem for ordinary differential equation in Menger probabilistic metric space

Abdolrahman Razani



AN EXISTENCE THEOREM FOR AN ORDINARY DIFFERENTIAL EQUATION IN Menger PROBABILISTIC METRIC SPACE

ABDOLRAHMAN RAZANI

Received 30 November, 2012

Abstract. In this article an existence theorem for an ordinary differential equation in probabilistic metric space is proved.

2010 Mathematics Subject Classification: 47H10; 54H25

Keywords: contractive map, fixed point, Menger probabilistic metric space, ordinary differential equations

1. INTRODUCTION

It is well known that the theory of probabilistic metric space is a new frontier branch between probability theory and functional analysis and has an important background, which contains the common metric space as a special case. Chang et al. [2] generalized contraction mapping principle in probabilistic metric space and as an application of this generalization, an existence theorem of solution of some kind of ordinary differential equations in probabilistic metric space is proved. Here, we consider an ordinary differential equations in the Menger probabilistic metric space, then we prove the existence of solution for this problem. Due to do this and for the sake of convenience, some definitions and notations are recalled from [1, 3–9].

Definition 1. A mapping $F : R \rightarrow R^+$ is called a *distribution function* if it is nondecreasing and left-continuous and it has the following properties:

- (1) $\inf_{t \in R} F(t) = 0$,
- (2) $\sup_{t \in R} F(t) = 1$.

Let D^+ be the set of all distribution functions F such that $F(0) = 0$ (F is a non-decreasing, left-continuous mapping from R into $[0, 1]$ such that $\sup_{t \in R} F(t) = 1$). Also denote by H the distribution function

$$H(t) = \begin{cases} 1, & t > 0, \\ 0, & t \leq 0. \end{cases}$$

Definition 2. A *probabilistic metric space* (briefly, *PM-space*) is an ordered pair (S, F) where S is a nonempty set and $F : S \times S \rightarrow D^+$ ($F(p, q)$ is denoted by $F_{p,q}$ for every $(p, q) \in S \times S$) satisfies the following conditions:

- (1) $F_{p,q}(t) = 1$ for all $t > 0$ if and only if $p = q$ ($p, q \in S$).
- (2) $F_{p,q}(t) = F_{q,p}(t)$ for all $p, q \in S$ and $t \in R$.
- (3) If $F_{p,q}(t_1) = 1$ and $F_{q,r}(t_2) = 1$ then $F_{p,r}(t_1 + t_2) = 1$ for $p, q, r \in S$ and $t_1, t_2 \in R^+$.

If only (2) and (3) hold, the ordered pair (S, F) is known as a *probabilistic semimetric space*.

Definition 3. A mapping $\Delta : [0, 1] \times [0, 1] \rightarrow [0, 1]$ is called a *triangular norm* (abbreviated, *t-norm*) if the following conditions are satisfied:

- (i) $\Delta(a, 1) = a$ for every $a \in [0, 1]$,
- (ii) $\Delta(a, b) = \Delta(b, a)$ for every $a, b \in [0, 1]$,
- (iii) $a \geq b, c \geq d \rightarrow \Delta(a, c) \geq \Delta(b, d)$ ($a, b, c, d \in [0, 1]$),
- (iv) $\Delta(a, \Delta(b, c)) = \Delta(\Delta(a, b), c)$ ($a, b, c \in [0, 1]$)

Definition 4. A *Menger probabilistic metric space* (briefly, *Menger PM-space*) (see [6]) is a triple (S, F, Δ) , where (S, F) is a probabilistic metric space, Δ is a t -norm and the following inequality holds:

$$F_{p,q}(t_1 + t_2) \geq \Delta(F_{p,r}(t_1), F_{r,q}(t_2)), \quad (1.1)$$

for all $p, q, r \in S$ and every $t_1 > 0, t_2 > 0$.

Definition 5. A triple (S, F, Δ) is called a *Menger probabilistic normed space* (briefly, *Menger PN-space*) (see [2]) if S is a real vector space, F is a mapping from S into D (for $x \in S$, the distribution function $F(x)$ is denoted by F_x and $F_x(t)$ is the value of F_x and $t \in R$) and Δ is a t -norm satisfying the following conditions:

- (i) $E_x(0) = 0$,
- (ii) $F_x(t) = H(t)$ for all $t > 0$ if and only if $x = 0$,
- (iii) $F_{\alpha x}(t) = F_x(\frac{t}{|\alpha|})$ for all $\alpha \in R, \alpha \neq 0$,
- (iv) $F_{x+y}(t_1 + t_2) \geq \Delta(F_x(t_1), F_y(t_2))$ for all $x, y \in E$ and $t_1, t_2 \in R^+$.

Schweizer, Sklar and Thorp [7] proved that if (S, F, Δ) is a Menger *PM-space* with $\sup_{0 < t < 1} \Delta(t, t) = 1$, then (S, F, Δ) is a Hausdorff topological space in the topology τ induced by the family of (ϵ, λ) -neighborhoods

$$\{U_p(\epsilon, \lambda) : p \in S, \epsilon > 0, \lambda > 0\},$$

where

$$U_p(\epsilon, \lambda) = \{u \in S : F_{u,p}(\epsilon) > 1 - \lambda\}.$$

Definition 6. Let (S, F, Δ) be a Menger PM -space with $\sup_{0 < t < 1} \Delta(t, t) = 1$.

- (1) A sequence $\{u_n\}$ in S is said to be τ -convergent to $u \in S$ (we write $u_n \rightarrow u$) if for any given $\epsilon > 0$ and $\lambda > 0$, there exists a positive integer $N = N(\epsilon, \lambda)$ such that $F_{u_n, u}(\epsilon) > 1 - \lambda$ whenever $n \geq N$.
- (2) A sequence $\{u_n\}$ in S is called a τ -Cauchy sequence if for any $\epsilon > 0$ and $\lambda > 0$, there exists a positive integer $N = N(\epsilon, \lambda)$ such that $F_{u_n, u_m}(\epsilon) > 1 - \lambda$, whenever $n, m \geq N$.
- (3) A Menger PM -space (S, F, Δ) is said to be τ -complete if each τ -Cauchy sequence in S is τ -convergent to some point in S .

Finally, we state a fixed point theorem form [2]. This theorem prove the existence of a fixed point in a Menger PM -space.

Theorem 1. Let (S, F, Δ) be a complete Menger PM -space with a t -norm Δ satisfying $\Delta(t, t) \geq t$ for all $t \in [0, 1]$. Let $T : S \rightarrow S$ be a mapping satisfying the following conditions:

$$F_{Tx, Ty}(t) \geq F_{x, y}\left(\frac{t}{k(\alpha, \beta)}\right),$$

for all $x, y \in S, t \geq 0$ and $\alpha, \beta \in (0, +\infty)$ with $F_{x, y}(\alpha) > 0$ and $F_{x, y}(\beta) < 1$, where $k(\alpha, \beta) : (0, +\infty)^2 \rightarrow (0, 1)$ is a function. Then T has exactly one fixed point in S .

In the next section, the existence of a solution for an ordinary differential equation in the Menger PN -space is proved.

2. AN EXISTENCE THEOREM

In this section we prove some lemmas to obtain the existence of solution for an ordinary differential equation in the Menger PN -space.

Lemma 1. If (R, F, Δ) is a Menger PN -space, then

$$|x| < |y| \implies F_x(t) > F_y(t),$$

for all $x, y \in R$ and $t \geq 0$.

Proof. Note that

$$F_x(t) = F_{\frac{x}{|y|}y}(t) = F_y\left(\frac{t}{|\frac{x}{y}|}\right) \geq F_y(t).$$

The last inequality holds, because $F_x(\cdot)$ is a nondecreasing function. \square

Definition 7. Let (R, F, Δ) be a Menger PN -space and $(E, \|\cdot\|_E)$ be a normed real vector space, we define a mapping $\tilde{F} : E \rightarrow D^+$ by

$$\tilde{F}_x(t) = F_{\|x\|_E}(t)$$

and denote the distribution function by \tilde{F}_x and the value of \tilde{F}_x at $t \in R$ by $\tilde{F}_x(t)$.

Proposition 1. *Let (R, F, Δ) be a Menger PN-space, then (E, \tilde{F}, Δ) is also a Menger PN-space.*

Proof. Note that

$$\tilde{F}_x(t) = F_{\|x\|_E}(t) \in D^+.$$

Secondly, $\tilde{F}_x(t)$ satisfies the next four conditions of the Definition 5.

- (i) $\tilde{F}_x(0) = F_{\|x\|_E}(0) = 0$,
- (ii) when $t > 0$, $F_x(t) = 1 \iff F_{\|x\|_E}(t) = 1 \iff \|x\|_E = 0 \iff x = 0$,
- (iii) $\tilde{F}_{\alpha x}(t) = F_{\|\alpha x\|_E} = F_{|\alpha|\|x\|_E}(t) = F_{\|x\|_E}(\frac{t}{|\alpha|}) = \tilde{F}_x(\frac{t}{|\alpha|})$,

and finally, by Lemma 1

$$\begin{aligned} (iv) \quad \tilde{F}_{x+y}(t_1 + t_2) &= F_{\|x+y\|_E}(t_1 + t_2) \geq F_{\|x\|_E + \|y\|_E}(t_1 + t_2) \\ &\geq \Delta(F_{\|x\|_E}(t_1), F_{\|y\|_E}(t_2)) \\ &= \Delta(\tilde{F}_x(t_1), \tilde{F}_y(t_2)). \end{aligned}$$

This ends the proof. \square

Proposition 2. *Let (R, F, Δ) be a Menger PN-space, then (R, G, Δ) is a PM-space, where*

$$G_{x,y}(t) = F_{x-y}(t)$$

Proposition 3. *Let (R, F, Δ) be a complete Menger PN-space with a continuous t -norm Δ . Then $(C([0, T]), R, \tilde{F}, \Delta)$ is also a complete Menger PN-space.*

Now, we can state the existence theorem for the solution of an ordinary differential equation in the probabilistic metric space as follow:

Theorem 2. *Let (R, F, Δ) be a complete Menger PN-space with a t -norm Δ satisfying $\Delta(t, t) \geq t$ for all $t \in [0, 1]$. Let f be Lipschitz mapping from $R \times \overline{U_{x_0}(\epsilon, \lambda)}$ into R . Then the following differential equation*

$$\begin{cases} \dot{x}(t) = f(t, x(t)), \\ x(0) = x_0. \end{cases}$$

has a solution in $C([0, \delta_0], R)$, where δ_0 is small enough.

Proof. Note that

$$\begin{cases} \dot{x}(t) = f(t, x(t)), \\ x(0) = x_0, \end{cases}$$

is equivalent to the following integral equation

$$x(t) = x_0 + \int_0^t f(s, x(s))ds.$$

Let $C([0, \delta_0], R)$ be the same as above, and

$$Tx(t) = x_0 + \int_0^t f(s, x(s))ds,$$

for all $x(.) \in C([0, \delta_0], \overline{U_{x_0}(\epsilon, \lambda)})$. Note that T is continuous and $T : C([0, \delta_0], \overline{U_{x_0}(\epsilon, \lambda)}) \rightarrow C([0, \delta_0], \overline{U_{x_0}(\epsilon, \lambda)})$ is contraction. In order to do this, we have

$$\begin{aligned} \tilde{F}_{Tx-Ty}(t) &= \tilde{F}_{\int_0^t (f(s, x(s)) - f(s, y(s))) ds}(t) \\ &= F_{\|\int_0^t f(s, x(s)) - f(s, y(s)) ds\|_\infty}(t) \\ &\geq F_{\delta_0 k \|x-y\|_\infty}(t) = F_{\|x-y\|_\infty}(\frac{t}{\delta_0 K}) = \tilde{F}_{x(t)-y(t)}(\frac{t}{\delta_0 K}) \end{aligned}$$

It means that

$$G_{Tx, Ty}(t) \geq G_{x, y}(\frac{t}{\delta_0 K}).$$

In addition, Proposition 3 shows $(C([0, \delta_0], R), G, \Delta)$ is a complete Menger PM -space and T is contraction, so by Theorem 1, T has exactly one fixed point in $C([0, \delta_0], R)$, i.e. there exists a unique $x(.) \in C([0, \delta_0], R)$ such that

$$Tx(t) = x(t).$$

It means that

$$x(t) = x_0 + \int_0^t f(s, x(s)) ds.$$

Thus $x(t)$ is a solution for

$$\begin{cases} \dot{x}(t) = f(t, x(t)), \\ x(0) = x_0, \end{cases}$$

Moreover, this solution is unique. \square

Example 1. Consider

$$\begin{cases} \dot{x}(t) = t + x(t), \\ x(0) = x_0, \end{cases} \quad (2.1)$$

By Theorem 2 the ordinary differential equation (2.1) for any $\delta_0 < 1$ has a unique solution.

Example 2. Consider

$$\begin{cases} \dot{x}(t) = t + \sin x(t), \\ x(0) = x_0, \end{cases} \quad (2.2)$$

By Theorem 2 the ordinary differential equation (2.2) for any $\delta_0 < 1$ has a unique solution.

REFERENCES

- [1] A. Aghajani and A. Razani, "Some completeness theorems in the menger probabilistic metric space," *Acta Mathematicae Applicatae Sinica*, vol. 10, pp. 1–8, 2008.
- [2] S. Chang, B. S. Lee, Y. J. Cho, Y. Q. Chen, S. M. Kang, and J.-S. Jung, "Generalized contraction mapping principal and differential equations in probabilistic metric spaces," *Proc. Amer. Math. Soc.*, vol. 124, pp. 2367–2376, 1996.
- [3] A. Razani, "A fixed point theorem in the menger probabilistic metric space," *New Zealand J. Math.*, vol. 35, pp. 109–114, 2006.
- [4] A. Razani and K. Fouladgar, "Extension of contractive maps in the menger probabilistic metric space," *Chaos, Solitons and Fractals*, vol. 34, pp. 1724–1731, 2007.
- [5] A. Razani and M. Shirdaryazdi, "A common fixed point theorem of compatible maps in menger space," *Chaos, Solitons and Fractals*, vol. 37, pp. 26–34, 2007.
- [6] B. Schweizer and A. Sklar, "Statistical metric spaces," *Pacific J. Math.*, vol. 10, pp. 313–334, 1960.
- [7] B. Schweizer, A. Sklar, and E. Thorp, "The metrization of statistical metric spaces," *Pacific J. Math.*, vol. 10, pp. 673–675, 1960.
- [8] H. Sherwood, "On e -spaces and their relation to other class of probabilistic metric spaces," *J. London Math. Soc.*, vol. 44, pp. 441–448, 1969.
- [9] Z. Shisheng, C. Yuqing, and G. Jinli, "Ekeland's variational principle and caristi's fixed point theorem in probabilistic metric space," *Acta Mathematicae Applicatae Sinica*, vol. 7, pp. 217–228, 1991.

Author's address

Abdolrahman Razani

Imam Khomeini International University, Department of Mathematics, P.O.Box 34149-16818, Qazvin, Iran

E-mail address: razani@sci.ikiu.ac.ir